


# ~ Intravenous Therapy ~

## 1. Keep dripping smooth :

when getting out of bed · Please use drip stand · Keep the IV higher than the injection site (about 90 cm or more) · and keep the injection site low and relaxed · avoid blood reflux to the infusion tube causing obstruction.



Do not adjust the IV drip rate by yourself. If you adjust drops too fast, causing cardiopulmonary discomfort; or too slowly, causing blood to clot and block the infusion tube.



If the drip is empty · The infusion line returns blood or there is a large amount of air, etc. · please notify the nursing staff.



When the intravenous indwelling needle is placed · There will still be pain where the needle was inserted · after about 10~20 minutes · it will be less pain.



## 2. Prevents needle displacement or slippage at the injection site :

When you need to change clothes during the drip injection · Please notify the nursing staff for assistance.



During any activity · remember not to press · twisting or pulling on the drip line or attachment · do not bump · push · touch the injection site · to avoid slippage.



When the injection site is at the joint · do not bend this joint; the arm in which the IV catheter is placed · do not lift items hard.



Do not peel off the tape holding the drip or remove the drip yourself.



## 3. Infection prevention :

Teach patients how to observe the injection site for complications, such as redness, swelling, heat, and pain.



If the skin around the injection site becomes red, pale, cold, swollen, painful, the IV connector becomes loose, bleeding or leaking, the IV drip is not flowing smoothly or not dripping, blood flows back, the adhesive tape becomes loose, or there is abnormal secretion, please notify the nursing staff immediately.



After removing the intravenous catheter, pay attention to whether the injection site is red, swollen, hot, painful, suppurative or bleeding. If any of the above conditions occur, please notify the nursing staff immediately and do not apply ice or heat compresses on your own.



Keep the indwelling needle site clean and dry to reduce the chance of infection: During the injection and drip period, you can clean your body by sponging. If you get wet, ask a nurse for assistance.




4. During intravenous injection, if you experience any discomfort such as difficulty breathing or a rapid heartbeat, tell the nurse immediately so they can assist you.



5. Provide patients with appropriate tube health education materials to improve patient compliance and prevent tube self-extraction and slippage.



6. Educate patients on the effects, side effects, and precautions of drugs.



*Intravenous Therapy* (圖-英文版)的檢閱修訂記錄

| 日期        | 版本  | 說明  | 備註 |
|-----------|-----|-----|----|
| 115/01/02 | 新制訂 | 新制訂 |    |